



More light on the $2\nu_5$ Raman overtone of SF₆: Can a weak anisotropic spectrum be due to a strong transition anisotropy?

Submitted by Florent Rachet on Thu, 03/26/2015 - 15:18

Titre More light on the $2\nu_5$ Raman overtone of SF₆: Can a weak anisotropic spectrum be due to a strong transition anisotropy?

Type de publication Article de revue

Auteur Kremer, David [1], Rachet, Florent [2], Chrysos, Michel [3]

Editeur American Institute of Physics

Type Article scientifique dans une revue à comité de lecture

Année 2014

Langue Anglais

Date Sep-01-2015

Numéro 3

Pagination 034308

Volume 140

Titre de la revue The Journal of Chemical Physics

ISSN 0021-9606

Résumé en anglais

Long known as a fully polarized band with a near vanishing depolarization ratio [$\eta_s = 0.05$, W. Holzer and R. Ouillon, Chem. Phys. Lett. 24, 589 (1974)], the $2\nu_5$ Raman overtone of SF₆ has so far been considered as of having a prohibitively weak anisotropic spectrum [D. P. Shelton and L. Ulivi, J. Chem. Phys. 89, 149 (1988)]. Here, we report the first anisotropic spectrum of this overtone, at room temperature and for 13 gas densities ranging between 2 and 27 amagat. This spectrum is 10 times broader and 50 times weaker than the isotropic counterpart of the overtone [D. Kremer, F. Rachet, and M. Chrysos, J. Chem. Phys. 138, 174308 (2013)] and its profile much more sensitive to pressure effects than the profile of the isotropic spectrum. From our measurements an accurate value for the anisotropy matrix-element $|\langle 000020 | \Delta\alpha | 000000 \rangle|$ was derived and this value was found to be comparable to that of the mean-polarizability $((000020), \alpha^{-1}, (000000))$. Among other conclusions our study offers compelling evidence that, in Raman spectroscopy, highly polarized bands or tiny depolarization ratios are not necessarily incompatible with large polarizability anisotropy transition matrix-elements. Our findings and the way to analyze them suggest that new strategies should be developed on the basis of the complementarity inherent in independent incoherent Raman experiments that run with two different incident-beam polarizations, and on concerted efforts to ab initio calculate accurate data for first and second polarizability derivatives. Values for these derivatives are still rarities in the literature of SF₆.

URL de la notice <http://okina.univ-angers.fr/publications/ua9241> [4]

DOI 10.1063/1.4861047 [5]

Titre abrégé J. Chem. Phys.

Liens

- [1] <http://okina.univ-angers.fr/dkremer/publications>
- [2] <http://okina.univ-angers.fr/florent.rachet/publications>
- [3] <http://okina.univ-angers.fr/michel.chrysos/publications>
- [4] <http://okina.univ-angers.fr/publications/ua9241>
- [5] <http://dx.doi.org/10.1063/1.4861047>

Publié sur *Okina* (<http://okina.univ-angers.fr>)